

Biochemical investigation on alcoholic and non-alcoholic liver diseases

¹Harpreet Kambir ²Ajjal Singh

¹Assistant Professor, ²Research Scholar

^{1,2}Department of Botany & Environmental Science,
N.P.P. College of Science, Raipur, Chhattisgarh

Abstract

In the present study, patients with liver diseases either due to alcohol or without alcohol compared with a group of normal healthy persons. Heavy drinkers showed significantly lower body weight and percent body fat, and low BMI compared with other groups. The percentage of hemoglobin and total number of RBC were found to be significantly decreased, whereas mean corpuscular volume significantly increased in alcoholic liver disease. However, no significant variation in either of the groups tested was observed in case of polymorphonuclear cells, lymphocytes, eosinophil and ESR values. Alcohol-induced liver injury is linked to oxidative stress as observed by decreased level of reduced glutathione and increased level of thiobarbituric acid reactive substances. Alanine amino transferase (ALT) and aspartate amino transferase (AST) are present in high concentration in hepatocytes. These enzymes leak into the circulation when hepatocytes or their cell membranes are damaged. Moderately elevated level of ALP and high GGT values are good discriminator of alcoholic patients. All these parameters in combinations may be useful indicator for identification and determination of severity of alcoholic liver diseases.

Key Words: Alcohol abuse, non-alcoholic liver diseases, hematological and Biochemical parameters.

Introduction

Alcohol abuse is a major public health problem in Tamil Nadu. In comparison to national average of alcohol consumption (22%), the average value for Tamil Nadu is higher (62%). In addition to this, the relapse to alcohol abuse, after de-addiction, is also as high as 42% (Gyatso and Bagdas, 1998). Severity of liver damage is often correlated with the amount of alcohol consumption in patients with a history of heavy alcohol abuse. However, alcoholic liver disease (ALD) not only depends on the total amount of alcohol consumed; drinking patterns and type of alcoholic beverage ingested are also playing important role in the development of ALD (Bellentani *et al.*, 2000). Most

patients develop fatty liver, which reverses on withdrawal of alcohol and is unlikely to progress to liver cirrhosis. In the liver, the acetaldehyde produced by oxidation of ethanol interacts with lipids and proteins, generating free radicals and impairing protein function (Fickert and Zatloukal, 2000).

Alcoholism is a condition resulting from excessive drinking of beverages that contain alcohol. The major health risk of alcoholism includes liver disease, heart disease, pancreatitis, central nervous system disorders and certain forms of cancer. The liver is particularly vulnerable to disease related to heavy drinking, most commonly termed as alcoholic hepatitis or cirrhosis. The progression of alcoholic liver disease is characterized by steatosis, inflammation, necrosis and cirrhosis. When severe hepatitis occurs, death is the outcome (Felver, 1990). Chronic consumption of alcoholic beverages is a primary cause of liver injury (Tuma, 2002). Hence, an attempt has been made to evaluate the effect of chronic alcohol consumption on serum bilirubin, serum proteins, serum lipids and hepatic marker enzymes.

Clinicians commonly fail to screen patients, and thus fail to recognize or treat alcoholism appropriately (Moore *et al.*, 1989). Clinical history that may suggest alcohol abuse or alcohol dependence includes the pattern, type, and amount of alcohol ingested, as well as evidence of social or psychological consequences of alcohol abuse. These may be suggested by other injuries or past trauma, such as frequent falls, lacerations, burns, fractures, or emergency department visits (Prytz and Melin, 1999). Biochemical tests have been considered to be less sensitive than questionnaires in screening for alcohol abuse, but may be useful in identifying relapse (Aalto and Seppa, 2005). Various questionnaires have been used to detect alcohol dependence or abuse, and include the CAGE, the Michigan Alcoholism Screening Test, and the Alcohol Use Disorders Identification Test (Soderstrom *et al.*, 1997). A structured interview, using instruments such as the Lifetime Drinking History, is often used as a gold standard for quantifying lifetime alcohol consumption (Skinner and Sheu, 1982).

Toxic substances generated during the metabolism of alcohol in the liver may contribute to the development of alcoholic liver disease. These substances include highly reactive molecules that can destroy vital cell components through a process called oxidation. Alcoholic liver diseases may be caused by oxygen radicals such as superoxide and hydroxyl radicals, generated during the metabolism of ethanol by the microsomal oxidising system (Das and Vasudevan, 2005). In the present study we have investigated changes in biochemical parameters in normal person, patients with alcoholic liver injury (both moderate and heavy drinkers), and non-alcoholic liver diseases. These

parameters were compared with normal values obtained from normal healthy people. Biochemical parameters included common laboratory tests.

Materials and Methods

The subjects were participants of the study. The participants were alcohol drinkers who came to attend the psychiatric clinic at Muthupet hospitals, Thiruvarur District for rehabilitation. Among them all alcohol dependent patient and liver disease patients volunteered for this study. Based on the number of years of consumption of alcohol, by the patients, they were divided into two groups.

A total of 240 subjects were studied: 85 subjects (55 males and 30 females) with alcoholic liver disease (ALD); 95 patients (60 males and 35 females) with non-alcoholic liver disease (NALD) and 60 normal healthy presenters (35 males and 25 females) without any reported disease. Alcoholic patients were further classified as described by Paton (7) into: a) High alcohol intake group (ALD-H; those had been drinking more than 80% alcohol per day for at least three years), and b) Moderate alcohol intake group (ALD-M; those had been drinking less than 25% alcohol per day). All participants were within same age group, similar economic status, non-smokers and similar dietary habits.

Blood samples: Venous blood samples were collected from control, liver disease and alcoholic patients. The blood samples were centrifuged at 5000 rpm for 20 minutes to obtain a clear supernatant (serum) which was stored at 4°C until its use for further analysis.

Hemoglobin concentration was determined using cyanmeth reagent (van Kampen and Zijlstra, 1965) and other hematological parameters were measured. Urea in serum was estimated by the kinetic method (Tiffany *et al.*, 1972) and creatinine level was estimated by rate of change in absorbance using alkaline picrate (Larsen, 1972). Uric acid was estimated as described by Gochman and Schmitz (1971). The total bilirubin level of serum was measured by the method of Malloy and Evelyn, (1937). The serum proteins like total proteins, albumin were estimated by the method of Lowry *et al.*, (1951). The serum lipids like total cholesterol, HDL, LDL, VLDL were estimated by the method of Warnick *et al.*, (1985) and triglycerides by Rifai *et al.*, (1998) method. Serum phosphatase levels (alkaline phosphatase and acid phosphatase activity) was determined by the method as employed by Linhardt and Walter (1963). Activities of alanine aminotransferase and aspartate aminotransferase in serum were determined by the method as described elsewhere (Bergmeyer and Bernt, 1963). The γ -glutamyltransferase activity of serum was assayed by the method

of Gowelock (1988). Extent of lipid peroxidation (Sinnhuber *et al.*, 1958) and reduced glutathione content (Beutler *et al.*, 1963) were also determined. All chemicals used were of analytical grade. Results have been expressed as mean \pm SEM (standard error). Statistical significance was determined by Student's 't' test for unpaired data. The values of significance were evaluated with 'p' values. The difference were considered significant at $p < 0.05$.

Results and Discussion

In the present study we examined the hematological, biochemical and enzymological changes in patients with liver diseases either due to alcohol or without alcohol were compared with a group of normal healthy persons. Alcohol has a variety of pathologic effects on hematopoiesis. It directly damages erythroid precursors, thereby contributing to macrocytosis and the anemic state of chronic alcoholics. Ethanol induces sideroblastic anemia, perhaps by direct interference with heme synthesis. Further, chronic ingestion of alcohol can lead to various types of hemolytic anemia caused by alterations in the erythrocyte membrane lipids that occur in association with alcoholic liver disease (Nordmann and Rouach, 1996). In the present study (Table 1), the percentage of hemoglobin and total number of RBC and WBC were found to be significantly decreased, whereas mean corpuscular volume (MCV) significantly increased in alcoholic liver disease with heavy alcohol intake in comparison to other groups. Because the red blood cell survive for 120 days after it has been released into the circulation, an MCV result may remain elevated for up to 3 months after a person has stopped drinking. But increase in MCV has been reported in other conditions such as thyroid disease, folate deficiency, recent blood loss and a number of hematological conditions, and liver disease from other causes (Whitfield *et al.*, 1978). Thus it cannot be taken as a sole parameter for alcoholic liver disease. However, no significant variation in either of the groups tested was observed in case of polymorphonuclear cells, lymphocytes, eosinophil and ESR values.

In the present study significant decrease in total protein was observed in all the tested groups when compared with normal healthy group, and heavy drinkers were found significantly low value compared to any other groups. Ethanol consumption slows down the rate of hepatic protein catabolism. Usharani *et al.*, (2012) reported that the effect of chronic alcohol consumption on serum bilirubin and total proteins. In our present study patients who had consumed alcohol for many years and heavy drinks exhibited hypoalbuminemia and hyperglobulinemia and the reversal of A/G ratio. Hypoalbuminemia is often associated with ascites and expansion of the

extravascular albumin pool at the expense of the intravascular albumin level (Marchesini *et al.*, 1991). In chronic alcoholism the liver cells are damaged and since albumin is solely synthesized in liver the concentration of serum albumin falls markedly in advanced parenchymal liver disease, when liver is damaged as that encountered in chronic alcohol consumers, the serum albumin levels are grossly decreased and the serum globulins are increased, thus leading to A/G ratio reversal (Chatterjea and Rana shinde, 1998). Thus the severity of hypoalbuminemia in alcoholics can serve as a criterion to assess the degree of damage to the liver cells. Das and Vasudevan (2005) studied decreased in total protein was observed in all the tested groups when compared with normal healthy group, and heavy drinkers were found significantly low value compared to any other groups. Ethanol consumption slows down the rate of hepatic protein catabolism.

The urea level was found to be within normal range in all the tested groups (Table 2). Though the increases in creatinine level of non-alcoholic liver disease group was found to be statistically significant in comparison to other groups, but no significant change was observed in alcoholic liver disease groups when compared with normal healthy persons (Table 3). Significant increase in uric acid level was observed in alcoholic liver disease groups with respect to other two groups, and high alcohol intake group showed significantly higher uric acid value when compared with moderate alcohol intake [ALD-M] group. Therefore, the bilirubin level in association with urea, creatinine, and uric acid may be used as markers in combination for ALD.

Liver plays an important role in lipid metabolism for two main reasons namely (a) The bile salts are formed in the liver which are necessary for emulsification and absorption of fats are excreted by liver (b) It is concerned with the metabolism of cholesterol. Marked alterations in lipid metabolism have been reported on chronic ethanol feeding (Usharani *et al.*, 2012). The accumulation of fat in the liver on chronic alcohol intake acts as a stimulus for the secretion of lipoproteins into the blood stream and the development of hyperlipidemia (Schapi *et al.*, 1965). Previous studies in our lab have shown that serum and tissue cholesterol levels increased with alcohol consumption in experimental rats (Senthilkumar *et al.*, 2003). Results of our present study in alcoholics also agree with the above findings.

Decreased fatty acid oxidation in the liver or increased fatty acid synthesis or both would increase the availability of substrate for lipoprotein synthesis. Such increased availability of substrate in the liver itself has to be postulated, since we found accumulation of lipids in the blood of alcoholic patients, without evidence that ethanol enhances clearance of lipoproteins from

the liver or affects lipid absorption or peripheral utilization (Parthasarathy *et al.*, 1989). Moreover, lipoproteins are chemically modified by oxidation. These oxidized or modified lipoproteins do not react with LDL receptors leading to esterification of cholesterol and conversion of macrophages to foam cells, thereby contributing to the hyperlipidemia observed on alcohol consumption. Cigarette smokers have high lipid peroxidation, which leads to damage the biomembrane and increased the plasma enzymes that reflect the protein concentration in plasma. Ramamurthy *et al.* (2012) have observed the increased levels of cholesterol, triglycerides, LDL and VLDL cholesterol where as HDL cholesterol was markedly decreased. The levels of cholesterol, triglycerides, LDL, VLDL-cholesterol levels were significantly increased but the levels of HDL-cholesterol were significantly decreased in all smokers as compared to non-smokers. Thus our study concludes that smokers have higher risk than that of non-smokers.

To find out the effects of alcohol consumption on hepatocellular injury the activities of AST and ALT levels in alcoholic patients were estimated (Table 3). Usharani *et al.*, (2012) studied showed no significant changes in AST and ALT levels in patients consuming alcohol of 6-15 years, while in patients consuming alcohol for a prolonged period, showed significantly elevated activities of AST and ALT as compared to those of the control subjects. Our results correlate with those of the previous findings which show elevated AST and ALT activities on alcohol consumption (Vajro *et al.*, 1992). Our data here suggest that the elevation was comparatively more in patients consuming alcohol for a prolonged period of more than 15 years.

SGOT and SGPT are present in high concentrations in cells of Heart, Liver etc. alcohol drinkers propagates the lipid peroxidation, which damage the biological membrane of the liver and heart. The enzymes are leaked out into blood and increased the level of SGOT and SGPT were observed in alcoholics in comparison to healthy controls as well as in heavy drinkers in comparison to moderate drinkers (Table 3). In our study, we have observed the SGOT and SGPT activities were significantly increased. Ramamurthy *et al.*, (2012) observed the enzymes content of the smokers showed a sharp decrease with respect to the non-smokers. On inhalation it passes through the lungs into the blood stream. It actually reacts with the hemoglobin of the red blood corpuscles forming a stable coordinated complex.

Gamma glutamyl transpeptidase (γ GT) is an enzyme produced in the bile duct and may be elevated in the serum of patients with bile duct diseases. Measurement of γ GT is an extremely sensitive test. It is induced by alcohol and its serum activity may be increased in heavy drinkers even in the absence of

liver damage or inflammation. In this study the serum γ GT levels were markedly increased in alcoholic patients, which was directly proportional to the number of years of consumption (Whitehead *et al.*, 1978). The elevation of γ GT alone with no other liver function test abnormalities often results from induction by alcohol. γ GT is often elevated in people who consume three or more alcoholic drinks (80% of ethanol or more) per day.

Significant increase in alkaline phosphatase, γ -glutamyltransferase, aspartate aminotransferase and alanine aminotransferase activities were observed in alcoholics in comparison to healthy controls as well as in heavy drinkers in comparison to moderate drinkers (Table 3). A similar change was also noted in case of ratio of aspartate aminotransferase and alanine aminotransferase. Alanine amino transferase (ALT) and aspartate amino transferase (AST) are present in high concentration in hepatocytes. These enzymes leak into the circulation when hepatocytes or their cell membranes are damaged. Although these aminotransferases are sensitive indicators of liver cell damage, neither alone is an ideal marker. In the non-alcoholic liver disease, the activities of AST, and ALT were increased significantly in comparison to normal healthy individuals. Though significant increase in activities of these enzymes were also observed in the moderate and heavy alcohol intake groups in comparison to normal healthy individuals, the rise was significantly less than the non-alcoholic liver disease. Serum aminotransferase concentrations are moderately raised in chronic and milder cases of acute viral or drug-induced hepatitis, autoimmune hepatitis, and alcoholic liver disease. Slightly raised serum aminotransferase levels characterize cirrhosis, non-alcoholic steato hepatitis, cholestatic liver disease, fatty liver and hepatic neoplasms (Schimdt and Schimdt, 1979).

Lactate dehydrogenase (LDH) is an intracellular enzyme particularly present in the kidney, heart, skeletal muscle, brain, liver and lungs. Increased LDH levels are usually found due to cell death and /or leakage from the cell of these organs Usharani *et al.*, (2012). In our present study also increased in the serum LDH concentration in alcoholic patients was found to be less significant than those of the AST / ALT levels in this study. Our results here show that the degree of elevation of all the hepatic marker enzymes can be used as biomarkers to evaluate approximately the number of years and dosage of alcohol consumed.

Conclusion

From the present study, it is very difficult to distinguish between the patients suffering from liver diseases either due to non-alcohol or moderate

alcohol intake. But alcoholic patients who were consuming high amount of alcohol are suffering from severe liver damage. Oxidative stress related enzymes and non-enzymes responded in the same way after liver is damaged. None of these parameters can be utilized as a marker for alcoholic liver disease. Antioxidants and stress related enzymes might be able to determine the degree of liver damage. Body weight, percent body fat, body mass index and hematological parameters are affected. Hyperbilirubinemia, hyperuricemia, hypoalbuminemia, high erythrocyte mean corpuscular volume, and normal urea and creatinine levels are common features of alcoholics. Monitoring γ GT, ALP, AST and ALT in combination is a sensitive means of detecting severity of alcohol induced liver damage. All the tests done in this study has found use in the early identification, and may help to monitor abstinence and relapse in response to outpatient treatment. The dosage of alcohol consumption which leads to coronary mortality can be done in future, so that the number of deaths due to alcoholism can be reduced. The differences between the groups might be based on the type of liver pathological condition rather than its etiology (i.e. alcohol and non-alcohol related causes).

References

1. Aalto, M., Seppa, K. 2005. Use of laboratory markers and the audit questionnaire by primary care physicians to detect alcohol abuse by patients. *Alcohol Alcohol*, 40: 520 – 523 .
2. Bellentani, S., Saccocio, G., Masutti, F., Giacca, M., Miglioli, L., Monzoni, A., Tiribelli, C. 2000. Risk factors for alcoholic liver disease. *Addiction Biology*, 5(3), 261-268.
3. Bergmeyer HU and Bernt E. 1963. Glutamate oxaloacetate transaminase; Glutamate pyruvate transaminase. In: *Methods of Enzymatic Analysis* (Ed. Bergmeyer HU) Academic Press, New York, pp.837-853.
4. Beutler E, Duron O and Kelly BM. 1963. Improved method for determination of blood glutathione. *J Lab Clin Med*, 61, 882-888.
5. Chatterjee MN and Rana Shinde MN. 1998. *Text book of Medical Biochemistry- 5th edition*. Jaypee brother medical publication (p) limited.
6. Das S.K. and D.M. Vasudevan. 2005. Monitoring Oxidative Stress in Patients with Non-Alcoholic and Alcoholic Liver Diseases. *Indian Journal of Clinical Biochemistry*, 20 (2) 24-28
7. Felver, ME, Merzey, E and Herlong, HF. 1990. Plasma tumor necrosis factor, a predicts decreased long term survival in severe alcoholic hepatitis. *Alcohol Clin. Exp Res*.31:117-134.

8. Fickert, P., Zatloukal, K., 2000. Pathogenesis of alcoholic liver disease. In: *Handbook of Alcoholism* (Eds. G. Zernig, A. Saria, M. Kurz, and S.S. O'Malley) Boca Raton, FL: CRC Press, 317- 323.
9. Gochman N and Schmitz JM. 1971. Automated determination of uric acid with use of a uricase-peroxidase system. *Clin Chem*, 17, 1154.
10. Gowelock, A.H. 1988. In: *Varley's Practical Clinical Biochemistry*. 6th edn. Heinemann Professional Publishing, p.519
11. Gyatso, TR and Bagdas, BB. 1998. In: *Health Status In Sikkim*. (Dept. of Health and Family Welfare, Govt. of Sikkim).
12. Larsen K. 1972. Creatinine assay by a reaction kinetic principle. *Clin Chem Acta*, 41, 209.
13. Linhardt K and Walter K. 1963. Phosphatase. In: *Methods of Enzymatic Analysis* (Ed. Bergmeyer HU) Academic Press, New York, p.799.
14. Lowry OH, Rose Borough NJ, Fsrer AL and Randall RJ. 1951. Protein measurement with Folin Phenol reagent. *J Biol Chem*. 193:265- 275.
15. Malloy HT and Evelyn KA. 1937. The determination of bilirubin with the photoelectric colorimeter. *J. Biol Chem*. 119:481- 490.
16. Marchesini G, Dioguardi FS and Bianchi GP. 1991. Long-term oral branched-chain amino acid treatment in chronic hepatic encephalopathy. A randomized double-blind casein-controlled trial. The Italian Multicenter Study Group. *J Hepatol*. 11:92-101.
17. Moore R.D, Bone LR and Geller. G. 1989. Prevalence, detection, and treatment of alcoholism in hospitalized patients. *JAMA*; 26: 403 - 407.
18. Nordmann R, Rouach H. 1996. Alcohol and free radicals: from basic research to clinical prospects. *Gastroenterol Hepatol*, 32(3), 128-133.
19. Parthasarathy S, Quinn MT and Schaenke OC. 1989. Oxidation role in monocyte recruitment and foam cell formation. *Atherosclerosis*. 1989;9:393-399.
20. Prytz, H and Melin, T. 1999. Identification of alcoholic liver disease or hidden alcohol abuse in patients with elevated liver enzymes. *J. Intern Med*, 233: 21- 26.
21. Ramamurthy, V., S. Raveendran, S.Thirumeni and S. Krishnaveni. 2012. Biochemical changes of cigarette smokers and non-cigarette smokers . *International Journal of Advanced Life Sciences*. I (1): 68 – 72.
22. Rifai N, Bachorik PS and Alberts JJ. 1998. Lipids, Lipoproteins and apolipoproteins in Burtis Rothschild MA, Oratz M and Schreiber SS. 1988. Serum albumin. *Hepatology*. 8:385-401.

23. Schapi RO, Scheig RL, Drummey G.D, Mendelson JH and Isserbacher KJ. 1965. Effects for prolonged ethanol injection on the transport and metal of lipids in Man. *New Eng. J. Med.*, 272: 610 - 615.
24. Schimdt E and Schimdt FW. 1979. Enzyme diagnosis in diseases of the liver and biliary system. In: *Advances in Clinical Enzymology. Vol.I.* (Eds. ESchimdt, FW Schimdt, I Trautschold, R Friedel) Basel: Karger; pp. 232-292.
25. Senthilkumar R, Viswanathan P and Nalini N. 2003. Glycine modulates hepatic lipid accumulation in alcohol induced liver injury. *Pol. J. Pharmacol.*, 5: 603-611.
26. Sinnhuber RO, Yu TC, Yu TC. 1958. Characterization of the red pigment formed in the thiobarbituric acid determination of oxidative rancidity. *Food Res*, 23: 626-630.
27. Skinner, HA and Sheu, W.J. 1982. Reliability of alcohol use indices. *The Lifetime Drinking History and the MAST. J. Stud. Alcohol.*, 43: 1157 – 1170.
28. Soderstrom, CA, Smith, GS and Kufera, JA. 1997. The accuracy of the CAGE, the Brief Michigan Alcoholism Screening Test, and the Alcohol Use Disorders Identification Test in screening trauma center patients for alcoholism. *J. Trauma*, 43: 962 - 969.
29. Tiffany TO, Jansen JM and Burtis CA. 1972. Enzymatic kinetic rate and endpoint analysis of substrate by use of GEMSAEC fast analyzer. *Clin. Chem.*, 18: 829.
30. Tuma DJ. 2002. Serial review: Alcohol, oxidative stress and cell injury free. *Radi. Biol. Med.*, 32:303-308.
31. Usharani, B., R. Vennila and N. Nalini. 2012. Biochemical changes in Alcoholics – A case control study. *International Journal of Research in Pharmaceutical and Biomedical Sciences*. 3 (1): 201 – 205.
32. Vajro P, Lofrano MM, Fontanella A and Fortunato G. 1992. Immunoglobulin complexed AST (“macro-AST”) in an asymptomatic child with persistent hypertansaminasemia. *J Pediatr Gastroenterol Nutr.*, 15:458-60.
33. van Kampen EJ and Zijlstra WG. 1965. Determination of hemoglobin and its derivatives. *Adv Clin Chem*, 8, 141-187.
34. Warnick GR, Nguyen T and Alberts AA. 1985. Comparison of improved precipitation methods for quantification of high density lipoprotein cholesterol. *Clin. Chem.*, 31:217.

35. Whitfield JB, Hensley WJ, Bryden D, Gallagher H. 1978. Some laboratory correlates of drinking habits. *Annals Clin Biochem*, 15: 297-303.